




Methicillin-Resistant *Staphylococcus* Information for Veterinarians



My Small Animal Patient has been diagnosed with a MRSA or MRSP, now what?

What is methicillin-resistant *Staphylococcus*?

- *Staphylococcus aureus* frequently colonize people, while *S. pseudintermedius* more commonly colonize dogs and cats.
- Methicillin-resistant *S. aureus* (MRSA) and *pseudintermedius* (MRSP) are defined by resistance to **oxacillin** and are resistant to all beta-lactams.
- MRSA and MRSP have public health importance because they can cause severe infections that are difficult to treat in people and pets.

Where did my patient get MRSA/MRSP?

- MRSA is most often spread to a pet from an infected or colonized person.
- MRSP is most often spread from another pet.
- MRSA/MRSP are typically spread through direct contact (hands, skin, nose, mouth, wounds) but can also spread via contaminated surfaces/equipment.
- Your patient might have already been colonized with MRSA/MRSP, then incurred a wound that became infected.
- MRSA and MRSP cause opportunistic infections in wounds, at surgical implants, and in the bladder.

Is there risk to my staff and other patients?

- With proper personal protective equipment and hand hygiene, the risk of spread is low.
- Immunosuppressed staff should avoid contact with MRSA/MRSP-infected patients when possible.
- Everyone can reduce spread by washing hands.
- Anyone concerned about exposure should consult with their healthcare provider.

How can we treat my patient?

- Treat infected pets with antibiotics based on culture and susceptibility.
- Topical mupirocin is often effective.
- Remove any nidus of infection (e.g., implants).
- Consult with a pharmacologist, microbiologist, or infectious disease specialist as needed for treatment guidance.
- Healthy pets who are colonized with MRSA/MRSP do not need to be treated.

General precautions

- Isolate hospitalized MRSA or MRSP patients and minimize number of staff caring for them.
- Wear gloves/gowns/masks when handling MRSA/MRSP patients, cleaning wounds, and changing bandages, and wash hands after all contact.
- Use caution not to cross-contaminate your hospital equipment and surfaces.
- Follow manufacturer guidelines for cleaning products (e.g., dilution, storage, contact time).

Where can I find more information?

- Kansas Veterinary Diagnostic Laboratory
(785) 532-5650 <http://www.ksvdl.org/>
- MU Veterinary Medical Diagnostic Laboratory
(573) 882-8367 <https://vmdl.missouri.edu/>
- [CDC MRSA Basics](#)

KANSAS STATE
UNIVERSITY

College of Veterinary Medicine

M College of
Veterinary Medicine
University of Missouri

This flier was originally developed by researchers at the Colleges of Veterinary Medicine at Kansas State University and the University of Missouri, and is funded in part by FDA Veterinary Laboratory Investigation and Response Network (U18FD006990).